SUPPLEMENTARY MATERIAL

Supplementary Table. Sequence of Events and Procedures.

Visit	Visit 1	Visit 2	Visit 3	Visit 4a	Visit 4b	Visit 5	Phone call
Day	-50 to -22	-21 ± 3	-14 ± 3	1	16 h (+1) from Visit 4a	8 ± 3	15 ± 3
PROCEDURE							
General Assessments							
Informed Consent and HIPAA ¹	Х						
Demographic Data	Χ						
Medical and Medication History	Х						
Update Medical and Medication		Х	Х	X	Х	Χ	Χ
History							
Allergic Skin Test	Χ						
Urine Pregnancy Test ²		Χ		X		Χ	
Review Inclusion/Exclusion	Х	Х	Х	Х			
Criteria							
Randomization				X			
AE (nTEAE/TEAE) Assessment	Χ	Χ	Χ	Χ	X ³	X_{6}	X
Allergen Challenge							
Titration CAC		Χ					
Confirmation CAC			Χ				
16 h Duration of Action CAC					Χ		
15 m Onset of Action CAC						Χ	
Signs and Symptoms		Х	Х	X ⁴	Χ	Χ	
Assessments							
Relief Drop Instillation ⁵		Χ	Χ		Χ	Χ	
Visual/Systems Exams							
Visual Acuity		Χ	Χ	Χ	Χ	X_{6}	
Slit Lamp Bio microcopy		Χ	Χ	Х	Χ	X ⁶	
Intraocular Pressure		Χ				Χ	
Dilated Fundoscopy		Х				Χ	
Investigational product (IP)							
IP Instillation				X ⁷		X_8	
Drop Comfort & Descriptor				Х			
Assessment							

¹ In the event that a subject has a medical condition, medication/contact lens washout, or needs to speak with the Investigator prior to Visit 1, the subject was given an informed consent form. Medical/medication history, demographics, skin test, and inclusion/exclusion review were performed at the time of informed consent signing prior to Visit 1 but must be confirmed at Visit 1 (with the exception of demographics and skin test).

doi: 10.18176/jiaci.0894

² Urine pregnancy testing was performed, if applicable, for females of childbearing potential.

³ Performed pre-CAC and post-CAC

⁴ Only pre-CAC

⁵ Relief medication could be administered to subjects at the end of Visits 2, 3, 4b, and 5, after all evaluations were completed.

⁶ Performed pre-CAC and post-CAC as part of the safety exit exam

⁷ 16 hour (+1 hour) pre-CAC

^{8 15 (+1)} minutes pre-CAC