

SUPPLEMENTARY MATERIAL

Table S1. Demographic and clinical features of the study population of 39 *Hymenoptera* venom-allergic patients and 22 controls.

Characteristics	Allergic (n=39)	Controls* (n=22)	P-value
Age in years (median, range)	46 (21-77)	35 (25-57)	0.0085
Female (%)	64.1	81.8	NA
Culprit (Nº)			
HBV	22	NA	NA
YJV	17		
Reaction severity	7/5/11/12/4	NA	NA
(Mueller grade; IV/III/II/I/LLR)			
ID skin tests (HBV/YJV) (% of positive)	100	ND	NA
Specific IgE (HBV/ YJV) (kU_A/L)	4.11 (1.96-12.00)	<0.35	<0.0001
Total IgE (HBV/ YJV) (kU_A/L)	49.80 (19.80-115.0)	21.40 (10.24-75.80)	0.14
Basal tryptase (kU_A/L)	4.89 (3.63-6.33)	2.80 (2.08-3.49)	<0.0001
BAT CD63 HBV/ YJV 1 µg/ml (%)	78.0 (59.20-86.70)	3.11 (1.20-4.85)	<0.0001

HBV: honey bee venom; YJV: yellow jacket venom; BAT: basophil activation test; ND: not determined.

*The controls had negative venom sIgEs and no history of an allergic reaction after being stung.

Table S2. Diagnostic performance of LAD2 MAT for *Hymenoptera* venom allergy.

Protocol	AUC	Diagnostic cut-off* [% CD63+ LAD2 cells]
StemPro-34 CM	0.58 (P=0.30)	3.5
StemPro-34 CM +IL33+IL6	0.62 (P=0.12)	5.1
Tyrode buffer+SCF+IL33+IL6	0.67 (P=0.03)	5.5

CM: complete media; AUC: area under the curve

*Determined by the Youden Index

Table S3. Comparison of demographic and clinical features between patients with nonsevere anaphylactic allergic reactions and patients with severe anaphylactic allergic reactions.

Feature	Nonsevere reaction*	Severe reaction*	P value
N (%)	27 (69.0)	12 (30.1)	
Female (%)	16 (59.3)	9 (75.0)	0.34
Age (y)	54.0 (35.3-63.0)	46.0 (31.0-60.0)	0.47
BST ($\mu\text{g/ml}$)	5.2 (4.0-6.9)	4.1 (3.7-5.8)	0.32
tIgE (kU/l)	46.4 (19.6-92.9)	96.7 (24.4-212.8)	0.11
sIgE to HBV/YJV (kU/l)	3.6 (1.6-10.6)	9.3 (2.3-36.3)	0.13
% CD63+ BAT anti-Fc ϵ RI 0.55 $\mu\text{g/ml}$	90.7 (71.9-94.8)	89.3 (82.2-94.4)	0.96
% CD63+ BAT HBV/YJV 1 $\mu\text{g/ml}$	76.7 (53.5-90.8)	79.1 (75.3-85.7)	0.65
% CD63+ BAT HBV/YJV 0.1 $\mu\text{g/ml}$	55.2 (36.1-73.1)	40.7 (18.9-78.8)	0.67
% CD63+ LAD2 MAT anti-IgE 10 $\mu\text{g/ml}$	19.1 (15.6-27.7)	25.0 (21.0-37.5)	0.07

BST: Basal serum tryptase; tIgE: total IgE; sIgE: specific IgE; BAT: basophil activation test; LAD2 MAT: LAD2 mast cell activation test.

Data are expressed as numbers (percentages) or medians (interquartile ranges).

*The severity was evaluated according to the Mueller scale. Large local, grade I and II reactions were considered nonsevere, and grade III and IV reactions were considered severe.

Table S4. Clinical and laboratory characteristics of 37 *Hymenoptera* venom allergic patients with nonresponding basophils

Characteristics	All (n=37)		
Age (median, range)	51 (24-80)		
Female (%)	22 (59%)		
Reaction severity	18/8/3/6/2		
(Mueller grade; IV/III/II/I/LLR)			
CD 63% basophils anti-FcεRI (median, IQR)	5.0 (3.5-6.9)		
CD63% LAD2 anti-IgE (median, IQR)	16.4 (12.9-21.1)		
Culprit	HBV (n=13)	YJV/hornet (n=15)	NI (n=9)
Skin tests HBV (N of positive)	9*/13	3/15	1*/9
Skin tests YJV (N of positive)	2*/13	8/15	2*/9
sIgE HBV (median, IQR)	5.45 (1.10-21.40)	<0.10 (<0.10-1.0)	0.50 (0.0-4.99)
sIgE YJV (median, IQR)	0.99 (0.30-1.62)	2.73 (1.44-9.37)	0.51 (<0.10-1.46)
LAD2 MAT HBV (median, IQR)	8.29 (5.11-13.10)	4.63 (3.26-5.72)	4.97 (3.34-9.72)
LAD2 MAT YJV (median, IQR)	1.76 (1.29-13.10)	3.26 (2.37-6.81)	2.85 (1.48-4.02)

HBV: honey bee venom; YJV: yellow jacket venom; NI: not identified; LAD2 MAT: LAD2 mast cell activation test.

*2 patients ND

Table S5. Detailed clinical and laboratory data of 37 patients with uninterpretable BAT results.

Patient ID	Culprit	Reaction severity (Mueller stage)	Skin tests		tIgE (kU/l)	sIgE (kU/l)		BAT (% CD63+)	LAD2 MAT (% CD63+)		
			HBV	YJV		HBV	YJV		HBV	YJV	anti-IgE
1	unknown	2	neg	neg	110	0.56	0.72	3.5	7.38	4.41	17.9
2	unknown	4	ND	ND	27.9	<0.1	0.51	2.6	2.4	2.0	23.1
3	YJV	4	neg	pos	27.5	<0.1	0.39	7.7	3.0	2.9	11.2
4	HBV	LLR	ND	ND	129	81.6	<0.1	4.6	17.0	1.55	15.2
5	unknown	4	neg	neg	81.1	8.5	<0.1	5.7	7.44	2.99	32.0
6	unknown	3	pos	neg	60.9	61.6	0.16	3.8	12.5	1.46	21.2
7	YJV	1	neg	neg	92.3	<0.1	1.44	5.3	4.24	1.62	15.7
8	HBV	3	pos	neg	140	30.1	12.6	3.1	12.1	1.19	20.9
9	HBV	4	pos	neg	22.1	11.4	0.22	9.0	5.88	0.77	11.2
10	HBV	3	pos	pos	50.5	0.93	1.37	4.8	8.29	4.53	17.3
11	hornet	4	neg	pos	38.2	<0.1	2.73	5.4	2.19	7.27	27.3
12	HBV	1	ND	ND	54.8	6.32	0.38	2.6	1.9	3.7	22.3
13	HBV	2	pos	neg	236	5.45	0.73	5.9	12.2	1.76	18.3
14	YJV	4	neg	neg	308	<0.1	1.57	6.6	4.6	3.1	16.4
15	HBV	1	pos	neg	81.7	2.88	11.3	5.0	12.3	5.9	19.0
16	HBV	4	pos	neg	176	12.7	1.48	5.8	17.3	11.2	24.1
17	YJV	3	neg	neg	65	0.452	2.56	3.5	9.95	7.28	25.0
18	hornet	4	pos	pos	203	1.11	2.90	4.1	2.4	2.1	8.95
19	HBV	4	pos	neg	26.3	3.96	1.76	8.3	13.9	1.39	16.6
20	HBV	3	pos	neg	16	1.03	0.57	2.8	8.18	2.97	13.7
21	HBV	4	pos	neg	181	36.3	0.991	4.7	7.34	1.83	18.5
22	YJV	4	neg	pos	28.4	<0.1	38.6	3.3	5.2	6.81	14.5
23	unknown	2	neg	pos	192	1.48	1.78	5.5	2.9	3.7	20.1
24	YJV	4	neg	pos	20.2	<0.1	1.33	9.0	2.5	0.9	3.83
25	YJV	4	neg	neg	313	<0.1	1.55	3.3	4.3	1.0	12.8
26	YJV	3	neg	neg	29.3	<0.1	0.36	3.2	2.1	3.0	12.8
27	HBV	1	neg	neg	45.2	1.17	<0.1	9.5	5.2	3.6	32.4
28	hornet	4	neg	pos	15.8	0.26	1.12	1.81	5.72	4.62	16.5
29	YJV	4	pos	pos	129	ND	31.7	9.7	4.2	4.4	22.9
30	unknown	4	neg	neg	42.9	0.5	<0.1	7.1	12.0	5.81	11.4
31	unknown	3	neg	neg	27.4	<0.1	1.14	5.2	4.4	1.5	8.51
32	hornet	4	neg	neg	91.7	<0.1	3.97	4.5	5.4	3.3	15.7
33	YJV	1	pos	pos	33.7	5.56	4.70	7.3	4.9	2.4	13.5
34	unknown	3	neg	pos	79.1	<0.1	5.0	5.0	5.0	3.6	13.4
35	unknown	LLR	ND	ND	117	0.48	0.42	7.1	3.8	2.9	12.0
36	hornet	4	neg	neg	41.6	1	9.37	4.1	6.15	6.85	12.9
37	hornet	1	neg	pos	109	37.9	2.98	6.0	23.0	2.46	14.4

tIgE: total IgE; sIgE: specific IgE; BAT: basophil activation test; LAD2 MAT: LAD2 mast cell activation test; LLR: large local reaction; HBV: honey bee venom; YJV: yellow jacket venom; ND: not determined.

Positive LAD2 MAT results are bolded.